



Murray Rosenblatt


1926–2019

BIOGRAPHICAL

Memoirs

*A Biographical Memoir by
Richard A. Davis*

©2021 National Academy of Sciences.
Any opinions expressed in this memoir
are those of the author and do not
necessarily reflect the views of the
National Academy of Sciences.




NATIONAL ACADEMY OF SCIENCES

MURRAY ROSENBLATT

September 7, 1926–October 9, 2019

Elected to the NAS, 1984

Murray Rosenblatt was a renowned researcher in both probability and statistics who is directly linked to the creation of several research seams that remain active today. Rosenblatt was the first to introduce kernel density estimation, now a standard nonparametric estimation technique that has generated significant new research. He co-authored, with Ulf Grenander, the acclaimed 1957 book *Statistical Analysis of Time Series*, a significant resource to both practitioners and researchers in time series for decades. And perhaps his most significant work is the introduction of the strong mixing condition, now one of the commonly used conditions to establish a central limit theorem (CLT).



By Richard A. Davis

Rosenblatt was born in New York City on September 7, 1926, the younger of two sons of Hyman and Ester (Goldberg) Rosenblatt, immigrants from Ukraine and Poland, respectively. Murray's older brother, David, and his spouse, Joan, were also mathematicians/statisticians of some note. Both had distinguished careers working in the federal government, with Joan spending her entire professional career at the National Institute of Standards and Technology (formally the National Bureau of Standards). Murray graduated from high school at the age of 16 and then proceeded to study mathematics at City College of New York (CCNY) in 1942. While at CCNY, Murray focused on courses in mathematical physics and thermodynamics. He was inspired by one of his analysis professors, Emil Post, who encouraged him to continue studying mathematics. After completing a bachelor of science degree at CCNY in 1946, he entered the graduate program in mathematics at Cornell University in 1946. Although having no previous experience in probability and statistics, he was exposed to two giants in the field—William Feller and Marc Kac—who in turn generated a great deal of synergy in probability that rubbed off on Murray. They were exciting times at Cornell, which attracted a

number of promising young visitors, including Joseph Doob, Monroe Donsker, Donald Darling, and Kai Lai Chung. Murray's first courses in probability theory and mathematical statistics were taught by Feller. In the end, he chose Kac as his Ph.D. supervisor because he found his personality more compatible with his own and because he allowed students more independence in their choice of research topics.

During his undergraduate years, Murray had a chance encounter with what was to be his future wife, Adylin Lipson, at the Fordham Road Library in the Bronx. Apparently, Adylin, known as Ady, did not make a positive first impression on Murray's mother. She blamed Ady for enticing Murray into walking home from the library in the rain, whereupon he developed a severe cold. Ady was also born and raised in the Bronx, received a bachelor's degree in social work from Hunter College in 1947 and would later earn a master's degree from San Diego State University in 1971. They were married in Ithaca in 1949, shortly after Murray received his Ph.D., and the couple spent one year in Ithaca before Murray began his academic career at the University of Chicago.

Career

After entering graduate school at Cornell, Murray's interest in analysis and thermodynamics naturally attracted him to Kac's work on statistical mechanics. He chose to work with Kac because he "sort of left you alone without saying you've got to do this or that so forth and so on. He let you go your own way." This advising philosophy would serve as Murray's mantra in mentoring his own Ph.D. students. As Murray described it, his dissertation, "On distributions of certain Wiener functionals" was "an attempt to mildly generalize some results of Kac" that were related to the seminal Feynman-Kac formula. The thesis, published in the *Transactions of the American Mathematics Society* under the title "On a class of Markov processes," derived properties of the Laplace transform of the distribution of certain integral functionals of Brownian motion. The resulting transform is found as a solution to a type of partial differential equation. Following his defense, Murray remained at Cornell for another year, supported by Kac's Office of Naval Research grant.

After finishing his work at Cornell, academic positions were scarce, and Murray was about to accept a government job when the University of Chicago stepped up and offered him a position in their statistics group. Murray believed that Kac likely played a behind-the-scenes role in making this connection. Murray joined the Committee of Statistics, the precursor to the Department of Statistics, at Chicago, that was then headed by W. Allen Wallis and included a number of budding young researchers that included

Jimmy Savage, Leo Goodman, Raghu Raj Bahadur, William Kruskal, and Charles Stein. It was at Chicago in 1951–52, when Murray first met Ulf Grenander, a young Swedish scholar. They would strike up a short but productive collaboration that produced a number of highly influential time series papers dealing primarily with estimation in the spectral domain. One of their few time domain papers considered estimation of a linear regression model with time series errors. They provided conditions on the regressors and the spectrum of the error process in order for ordinary least squares (OLS) estimates to be asymptotically efficient. This is one of the fundamental results in times series analysis. Much of their collaborative work appeared in their much-acclaimed 1957 book *Statistical Analysis of Time Series*. This book became a must-read for both practitioners and researchers in time series for decades. With the large number of bright young researchers and an active visitors program, the environment at Chicago was nearly ideal for someone like Murray. In addition to writing a couple of papers with visiting scholar Joe Hodges, he was also inspired by Bahadur. Murray credits his discussions with Bahadur at the beginning of his “little paper” on nonparametric density estimation,¹ which appeared in the *Annals of Mathematical Statistics* in 1956. Of course this “little paper,” which currently has around 5,000 citations, was the first to introduce kernel density estimation. This has now become one of the standard nonparametric estimation techniques and generated a rich and important line of research activity.

In 1956, Murray left Chicago for the mathematics department at Indiana University, where he struck up a collaboration with Julius Blum. The stint at Indiana lasted only a few years before he landed at Brown University in 1959. Returning to the East Coast had many attractions, including the opportunity to meet regularly with researchers at nearby Bell Laboratories. There he interacted with David Slepian and others who triggered his interest in modeling physical science phenomena. Motivated by a problem in human vision, Murray and Slepian wrote a paper on a n th order Markov process in which n variables are independent but $(n+1)$ variables are dependent.² While visiting Bell Labs, Murray met one of John Tukey’s Ph.D. students, David Brillinger, who was working on higher-order properties of times series models. This interaction developed into a series of collaborative papers on higher-order spectra.

During his Brown years, Murray had an opportunity to visit the Statistics Department at Columbia University. It was there that he wrote perhaps his most well-known paper, “A central limit theorem and a strong mixing condition,”³ which introduced the strong mixing condition. This work was motivated in part by an idea of Serge Bernstein that broke up sequences of random variables into nearly independent blocks of random

variables. The condition on the blocks, which specifies that they become roughly independent as the blocks grow in size and the separation increases, is in essence the strong mixing condition.

Murray left Brown for the newly created University of California at San Diego (UCSD) in 1964. Scripps Institution of Oceanography predated the university, and its strong research group that was analyzing a multitude of time series, not to mention the San Diego weather, was a major attraction. Murray arrived in 1964, one year before the first undergraduates were admitted. These were heady times to be part of the new faculty that in a short time would turn UCSD into a world-class research university. UCSD was fortunate to have attracted a number of talented young probabilists and continues to this day to have a major presence in probability theory.

Technical Contributions

Strong Mixing and the Central Limit Theorem

In his most celebrated paper, Rosenblatt introduced the notion of strong mixing for stationary time series. If $\{X_t\}$ is a stationary time series, then strong mixing specifies the rate at which events defined by two sets of random variables separated by m -time lags become independent. So for example, if $\mathcal{B}_0 = \sigma(X_s, s \leq 0)$ and $\mathcal{F}_n = \sigma(X_s, s \geq n)$, are σ -fields generated by the past relative to time 0 and the future of time n , then the time series is said to be strong mixing if

$$\sup_{A \in \mathcal{B}_0, B \in \mathcal{F}_n} |P(A \cap B) - P(A)P(B)| \leq \alpha(n),$$

where the mixing function $\alpha(\cdot)$ decreases to zero as $n \rightarrow \infty$. Although other mixing conditions existed around that same time, including m -dependence in which random variables separated by more than m -time lags are independent, Rosenblatt, in his understated manner, wrote, “The strong mixing condition used in this paper seems to be a more intuitively appealing formalization of this notion than most others.” Of course he was right, as the strong mixing condition became one of the most commonly used conditions to establish a central limit theorem (CLT). Under a mean zero and finite $2+\delta$ -moment condition on X_t , a growth restriction on the variance of the partial sums, $S_n = \sum_{t=1}^n X_t$, and a convergence rate on the mixing function to 0, Rosenblatt showed that S_n/\sqrt{n} was asymptotically normally distributed, although his results were more general than stated here. The idea of the proof was based on Bernstein’s big-block small-block construction. Essentially, the sequence of random variables X_1, \dots, X_n is separated

into k_n blocks of size $p_n + q_n$, where p_n and q_n are the sizes of the big and small blocks, respectively with $n = k_n(p_n + q_n)$. Now the big blocks of random variables are separated by q_n lags, and hence these blocks of random variables become independent via the strong mixing condition. The sum over the small blocks become negligible in the limit, and hence the sum S_n can be viewed as roughly a sum of k_n independent random variables that are asymptotically normal by the standard CLT.

Kernel Density Estimation

Rosenblatt's 1956 seminal paper on nonparametric density estimation introduces the kernel density estimate.⁴ The objective of this paper was to construct an estimate of the probability density function (pdf) $f(x)$ from a sample of independent and identically distributed random variables X_1, \dots, X_n having density f . The motivation for the kernel estimate comes from taking the difference of the empirical distribution function $F_n(x) = \frac{\#\{i \leq n: X_i \leq x\}}{n}$. So an obvious estimate of the pdf is given by the difference,

$$f_n(y) = \frac{F_n(y+h) - F_n(y-h)}{2h},$$

where $h=h_n$ is the bandwidth that is a function of the sample size n that converges to 0 with increasing n . Using well-known properties of F_n , it is straightforward to derive the corresponding statistical properties of f_n . Assuming the existence of three derivatives of f , he showed that the optimal choice of the bandwidth that minimizes the asymptotic integrated mean squared error, $\int_{-\infty}^{\infty} E|f_n(y) - f(y)|^2 dy$, is

$$h_n = kn^{-\frac{1}{5}}, \quad \text{where} \quad k = \left[\frac{9}{2 \int_{-\infty}^{\infty} |f'''(y)|^2 dy} \right]^{\frac{1}{5}}.$$

This naive estimate is an example of a kernel density estimate in which the kernel $w(x) = \frac{1}{2}$, for $|x| \leq 1$, and 0, otherwise. More generally, this paper considered kernel density estimates given by

$$f_n(y) = \int_{-\infty}^{\infty} w_n(y-u) dF_n(u) = \frac{1}{n} \sum_{t=1}^n w_n(y - X_t),$$

where $w_n(u) = \frac{1}{h} w\left(\frac{u}{h}\right)$, and $w \geq 0$ satisfies, $\int_{-\infty}^{\infty} w(u) du = 1$, $\int_{-\infty}^{\infty} uw(u) du = 0$, and $\int_{-\infty}^{\infty} |u|^3 w(u) du < \infty$. The mean-squared error of these estimates is of order $O(n^{-\frac{4}{5}})$. This basic idea was extended by many others to additional sorts of problems, including nonparametric regression estimates. A great deal of effort went into designing optimal

kernel functions. Rosenblatt continued his research on nonparametric density estimation throughout much of his research life, and one of his most noted papers is with Bickel on global measures of density estimates, which appeared in the *Annals of Statistics* in 1973.⁵ The idea was to consider the kernel density estimate a) for constructing uniform confidence bands for f of the form $\hat{f}_n \pm c_n \sqrt{\hat{f}_n}$ and b) for constructing goodness of fit tests for the null hypothesis $H: f=f_0$ using the test statistic

$$\max_x \frac{|f_n(x) - f_0(x)|}{\sqrt{f_0(x)}}.$$

The limit theory was based on the heuristic that the process $U_n(x) := \sqrt{nh_n} \frac{f_n(x) - f(x)}{\sqrt{f(x)}}$, which was asymptotically independent for distinct and fixed values of x , behaved locally like a stationary Gaussian process. The resulting test statistic could then be described as a maxima of a Gaussian process over an increasing interval on the real line. The results were ahead of their time, as the Bickel and Rosenblatt paper also made detailed use of the limiting distribution (Gumbel) of the maxima of a Gaussian process under different smoothness conditions.

Limit Theory for Long Memory Processes

In his 1960 Berkeley Symposium paper, "Independence and Dependence," Rosenblatt considered limit theory for partial sums of stationary sequences exhibiting long memory. In such a setting, the CLT he had established for strongly mixing sequences no longer applied. In this paper, he considers a function of a stationary Gaussian process $\{Z_t\}$ with mean 0 and autocovariance function $r(h) = (1 + h^2)^{-\frac{\gamma}{2}}$, with $0 < \gamma < \frac{1}{2}$. He chose the simplest nonlinear function of $\{Z_t\}$ (linear functions would again be Gaussian and not of interest) given by $\mathbf{X}_t = \mathbf{Z}_t^2 - 1$, which again is a mean zero stationary time series. But this process is not strong mixing. The spectral density function of the Z_t has a singularity of the form $|\omega|^{\gamma-1}$ in a neighborhood of the origin. Rosenblatt then established the limit distribution of the normalized partial sums, $n^{\gamma-1} \sum_{t=1}^n \mathbf{X}_t$. Given the long-memory of the process, the variance of the partial sums grows at a much slower rate and hence the normalization is much smaller, $n^{\gamma-1}$ compared to the usual $n^{-\frac{1}{2}}$. In addition the limit distribution is non-normal. This example has generated a great deal of interest, and it was picked up by others, specifically M. S. Taquq⁶ and R. L. Dobrushin and P. Major.⁷ They extended this notion to a wide range of processes and were able to classify functional limits of the partial sum process.

Bispectrum

Much of the early work in time series had a strong frequency domain flavor, with focus on the second-order spectrum. It was found that second-order models were not sufficiently rich to model a range of physical phenomena. In the late 1950s and early 1960s, A. Kolmogorov urged the young scholars V. P. Lennov and A. N. Shiryaev to consider higher-order models with a view towards describing nonlinear behavior. Such models require analysis that goes beyond the standard spectral analysis, which only dealt with second-order properties (covariance function) of the process. In the mid-1960s, Rosenblatt began to work in a similar vein on higher-order spectra. Around this time, a promising Princeton Ph.D. student, David Brillinger, who was working at Bell Labs, was inspired by Rosenblatt's work and an important collaboration was formed. In a series of influential papers in 1967, Brillinger and Rosenblatt developed the asymptotic theory for estimates of the cumulant spectrum under mixing conditions. This research set the stage for identification of nonlinear time series models. Rosenblatt continued working on higher-order spectral theory throughout his professional career. These ideas played a key role in modeling non-minimum phase and non-Gaussian linear processes. These ideas also played a role in his books *Stationary Sequences and Random Fields* and *Gaussian and Non-Gaussian Linear Time Series and Random Fields*.

Other Major Contributions

Rosenblatt made major contributions in a number of other areas as well. Much of this work seemed to be motivated by a conjecture by N. Wiener⁸ that gets at the heart of general time series modeling. The setup is the following. If $\{X_j\}$ is a stationary process with $\mathcal{B}_n = \sigma(X_j, j \leq n)$ then when does the process have a one-sided causal representation, i.e., when does X_n have the representation,

$$X_n = f(\xi_n, \xi_{n-1}, \dots),$$

for some iid sequence $\{\xi_t\}$. Wiener conjectured that a necessary and sufficient condition for such a representation is that the backward tail σ -field be trivial, $\mathcal{B}_{-\infty} = \bigcap_n \mathcal{B}_n = \{\emptyset, \Omega\}$. Rosenblatt⁹ showed this was true for Markov chains with a countable state space. Rosenblatt^{10,11} discussed when this conjecture holds and fails. Aside from the full scope of when Wiener's conjecture holds, one can see traces of this notion in many of Rosenblatt's research areas. This includes his interest on limits of convolutions of probability measures on compact semigroups and in the deconvolution problem. The basic setting of the deconvolution problem is that the observations X_t come from a linear system driven by iid noise, i.e.,

$$X_t = \sum_{j=-\infty}^{\infty} \psi_j \xi_{t-j},$$

where the $\{\psi_j\}$ represents a linear filter and $\{\xi_t\}$ is iid noise. Then the question is, when can one recover the noise sequence? If the filter is known, this is relatively straightforward. But if the filter is unknown, then the goal is to *deconvolve* the filter from the noise. In a case in which the noise is Gaussian, then there is an identification problem. Rosenblatt studied this problem in the non-Gaussian setting using higher-order spectra and likelihood methods when the distribution of the noise is known. It is of interest to know when the filter is minimum phase, which corresponds to the case that the representation is one sided, $X_t = \sum_{j=0}^{\infty} \psi_j \xi_{t-j}$, depending only on the past of the $\xi_s, s \leq t$. So Wiener's conjecture comes full circle to modern time-series modeling.

Finally, there is the three-page Rosenblatt paper, "Remarks on a multivariate transformation" in the *Annals of Mathematical Statistics*.¹² The rather straightforward transformation described in this paper maps a random vector $X=(X_1, \dots, X_k)$ with an absolutely continuous distribution into a random vector in which the components are iid uniform on $(0,1)$. This is the multivariate analogue of the standard one-dimensional probability integral transform. Although this is not a deep result, it has proven extremely handy in modern statistics, especially in the context of copula modeling and goodness of fit of multivariate distributions. This paper has nearly 3,000 citations.

After Murray became Distinguished Professor Emeritus at UCSD in 1994, he remained actively engaged in research until the last few years of his life. After a courageous fight with cancer, Ady passed away in 2009. The Rosenblatts had two children, daughter Karin (currently of Champaign, Illinois) and son Daniel (of Live Oak, Texas). Murray advised twenty-two dissertations, and to many of these Ph.D. students, he and Ady served as surrogate parents during their graduate school years. He was there for his students with great encouragement and support. In celebration of Murray's 90th birthday, the Murray and Adylin Rosenblatt Endowed Lecture Series in Applied Mathematics was created at UCSD in 2016.¹³

Concluding Remarks

Murray Rosenblatt had amazing intuition and insight about random phenomena—it was a sixth sense about how things should work. He was a scholar who seemed to never have forgotten a historical fact or theorem. To his Ph.D. students, he appeared to be the

most knowledgeable person, on topics from geography to history to the physical sciences and mathematics, that they ever encountered. While Murray had a soothing and relaxing style of lecturing, he could be laconic in both his writings and verbal communication. The latter could be especially perplexing to new researchers trying to decode his thoughts and insights on various mathematics problems. It was not uncommon for a former Ph.D. student or young colleague to only fully grasp what Rosenblatt had clearly understood and tried to convey years earlier: “Oh, that is what Murray meant” was a common thought. Murray’s advising approach replicated the one he experienced with Kac. He would let his students work on what they wanted and left them to their own devices. But he offered clear guidance, especially when students were struggling to find their way. He made all of his students feel important regardless of their aptitude and provided behind the scenes support in job searches and other endeavors to help further their careers.

Later in life Rosenblatt became fascinated with Chinese silk and became a quasi-expert, an expression he liked to use, on all things related to the manufacturing and production of the fabric. He loved going on long walks and hiking in the mountains whenever he had the chance. Even at an advanced age, it was difficult for his younger companions to keep up with his pace. In later years, Murray and Ady very much enjoyed spending holidays in Hawaii with his brother David and sister-in-law Joan.

Stigler’s law of eponymy states that no scientific discovery is named after its original discoverer. Murray would occasionally echo sentiments similar to this law to his Ph.D. students, who although they were enthusiastic, they were naïve and did not know better.¹⁴ So it is somewhat ironic that a number of terms in probability and statistics now bear the Rosenblatt name. This includes the Rosenblatt transformation and the Rosenblatt process, and sometimes strong mixing is referred to as Rosenblatt mixing. Murray also tended to eschew labels. Was he a probabilist, a statistician, a mathematician? He spent his entire academic life in a department of mathematics as opposed to a statistics or engineering department. His affinity towards mathematics was never more evident than during one of our last conversations when I asked, “What are you thinking about these days?” Murray responded, “I am thinking about the foundations of mathematics.” He then pointed me to the book on his desk, *Princeton Companion to Mathematics*, a heavy read that he was currently working through. Question answered—always a mathematician.

MAJOR HONORS

- 1955 Fellow, Institute of Mathematical Statistics
- 1965-66 Guggenheim Fellow
- 1970 Wald Lectures
- 1971-72 Guggenheim Fellow
- 1975 Fellow of the American Association for the Advancement of Science
- 1979 Overseas Fellow, Churchill College, Cambridge University
- 1984 Elected to the National Academy of Sciences
- 2009 Fellow of the Society for Industrial and Applied Mathematics
- 2014 Fellow of the American Mathematical Society

REFERENCES

1. Rosenblatt, M. 1956. Remarks on some nonparametric estimates of a density function. *Ann. Math. Statist.* 27:832–837.
2. Rosenblatt, M., and D. Slepian. 1962. Nth order Markov chains with every N variables independent. *J. Soc. Indust. Appl. Math.* 10:537–549.
3. Rosenblatt, M. 1956. A central limit theorem and a strong mixing condition. *Proc. Natl. Acad. Sci. U.S.A.* 42:43–47.
4. Rosenblatt, M. 1956. Remarks on some nonparametric estimates of a density function. *Ann. Math. Statist.* 27:832–837.
5. Bickel, P. J., and M. Rosenblatt. 1973. On some global measures of the deviations of density function estimates. *Ann. Stat.* 1(6):1071–1095.
6. Taqqu, M. S. 1975. Weak convergence to fractional brownian motion and to the Rosenblatt process. *Z. Wahrscheinlichkeitstheorie verw Gebiete* 31:287–302.
7. Dobrushin, R. L., and P. Major. 1979. Non-central limit theorems for non-linear functional of Gaussian fields. . *Wahrscheinlichkeitstheorie verw Gebiete* 50:27–52.
8. Wiener, N. 1958. *Nonlinear Problems in Random Theory*. Cambridge, Mass.: MIT Press.
9. Rosenblatt, M. 1960. Stationary Markov chains and independent random variables. *J. Math. Mech.* 9:945–950.
10. Rosenblatt, M. 2009. Stationary processes and a one-sided representation in terms of independent identically distributed random variables. In: *Dependence in Probability, Analysis and Number Theory*, ed. Herold Dehling, pp. 311-315. Heber City, Utah: Kendrick Press.
11. Rosenblatt, M. 2009. A comment on a conjecture of N. Wiener. *Stat. Probabil. Lett.* 79(3):347–348.
12. Rosenblatt, M. 1952. Remarks on a multivariate transformation. *Ann. Math. Statist.* 23:470–472.
13. Contributions in memory of Murray and Ady can be made to this lecture series via <https://giveto.ucsd.edu/giving/home/gift-referral/4810f5ce-6b34-498b-9477-9de96de2b82c>.
14. Brillinger, D. R., and R. A. Davis. 2009. A conversation with Murray Rosenblatt. *Stat. Sci.* 24(1):116–140.

SELECTED BIBLIOGRAPHY

- 1949 *On distributions of certain Wiener functionals*. Ph.D. thesis, Cornell University.
- 1951 On a class of Markov processes. *Trans. Amer. Math. Soc.* 71:120–135.
- 1952 Remarks on a multivariate transformation. *Ann. Math. Statist.* 23:470–472.
- 1953 With K. A. Brownlee and J. L. Hodges Jr. The up-and-down method with small samples. *J. Am. Stat. Assoc.* 48(262):262–277.
- With J. L. Hodges Jr. Recurrence-time moments in random walks. *Pacific J. Math.* 3:127–136.
- 1954 With U. Grenander. Regression analysis of time series with stationary residuals. *Proc. Natl. Acad. Sci. U.S.A.* 40(9):812.
- 1956 With R. Blum. A class of stationary processes and a central limit theorem. *Proc. Natl. Acad. Sci. U.S.A.* 42:412–413.
- A central limit theorem and a strong mixing condition. *Proc. Natl. Acad. Sci. U.S.A.* 42:43–47.
- Remarks on some nonparametric estimates of a density function. *Ann. Math. Statist.* 27:832–837.
- 1957 With U. Grenander. *Statistical Analysis of Time Series*. New York: Wiley.
- 1958 With C. J. Burke. A Markovian function of a Markov chain. *Ann. Math. Statist.* 29:1112–1122
- With N. Weiner. *Nonlinear Problems in Random Theory*. New York: Technology Press of Massachusetts Institute of Technology.
- With C. J. Burke. Consolidation of probability matrices. *Bull. Inst. Internat. Statist.* 36, no. 3, 7–8.
- 1959 Functions of a Markov process that are Markovian. *J. Math. Mech.* 8:585–596.
- Stationary processes as shifts of functions of independent random variables. *J. Math. Mech.* 8:665–682.

- 1960 Stationary Markov chains and independent random variables. *J. Math. Mech.* 9:945–950.
- Independence and dependence. In *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Vol. 1*. J. Neyman, ed. Pp. 431–443. Berkeley and Los Angeles: University of California Press.
- 1962 With D. Slepian. N th order Markov chains with every N variables independent. *J. Soc. Indust. Appl. Math.* 10:537–549.
- Random Processes*. Oxford, United Kingdom: Oxford University Press.
- 1967 With D. R. Brillinger. Computation and interpretation of k -th order spectra. In *Spectral Analysis of Time Series*. B. Harris, ed. Pp. 189–232. New York: Wiley.
- With D. R. Brillinger. Asymptotic theory of estimates of k -th order spectra. *Proc. Natl. Acad. Sci. U.S.A.* 57:206–210.
- 1971 *Markov Processes, Structure and Asymptotic Behavior: Structure and Asymptotic Behavior*. New York: Springer/Verlag.
- 1973 With P. J. Bickel. On some global measures of the deviations of density function estimates. *Ann. Stat.* 1(6):1071–1095.
- 1980 With K. S. Lii. Deconvolution and estimation of transfer function phase and coefficients for non-Gaussian linear processes. *Ann. Stat.* 10:1195–1208.
- 1997 Sun, T.C. Murray Rosenblatt: His contributions to probability and statistics. *J. Theor. Prob.* 10(2):279–286.
- 2000 *Gaussian and non-Gaussian linear time series and random fields*. Berlin/Heidelberg, Germany: Springer Science & Business Media.
- 2009 Stationary processes and a one-sided representation in terms of independent identically distributed random variables. In *Dependence in Probability, Analysis and Number Theory*, ed. Herold Dehling, pp. 311–315. Heber City, Utah: Kendrick Press.
- A comment on a conjecture of N. Wiener. *Stat. Probabil. Lett.* 79(3):347–348.
- 2011 Davis, R. A., K. S. Lii, and D. N. Politis, eds. 2011. *Selected Works of Murray Rosenblatt*. Berlin/Heidelberg, Germany: Springer Science & Business Media.

Published since 1877, *Biographical Memoirs* are brief biographies of deceased National Academy of Sciences members, written by those who knew them or their work. These biographies provide personal and scholarly views of America's most distinguished researchers and a biographical history of U.S. science. *Biographical Memoirs* are freely available online at www.nasonline.org/memoirs.